













Ineffective communication nullifies any attempt at care provision. The quality of care health workers provide does not matter if patients do not understand what they are being told. This leads to negative patient outcomes, increased utilization of emergency and inpatient services and a greater cost burden on healthcare systems. Effective communication includes health literacy, cultural competency and language barriers. Interventions to address each component should be incorporated into every level of healthcare organizations and fit seamlessly into preexisting workflows. Corresponding action plans should be created for each component in order to determine baseline capabilities and track improvements. Optimizing all components of effective communication will improve patient outcomes and lead to greater monetary savings which can then be reinvested in the organization.

-
1. BusinessDictionary.com. <http://www.businessdictionary.com/definition/effective-communication.html>. Accessed August 3, 2019. 
 2. Schyve PM. Language Differences as a Barrier to Quality and Safety in Healthcare: The Joint Commission Perspective. *J Gen Intern Med*. 2007;22(S2):360-361. 
 3. AHRQ Health Literacy Universal Precautions Toolkit. <https://www.ahrq.gov/professionals/quality-patient-safety/quality-resources/tools/literacy-toolkit/index.html>. Published February 20, 2015. Accessed September 4, 2017. 
 4. Welch VL, Vangeest JB, Caskey R. Time, Costs, and Clinical Utilization of Screening for Health Literacy: a Case Study Using the Newest Vital Sign (NVS) Instrument. *J Am Board Fam Med*. 2011;24(3):281-289. 
 5. Sudore RL, Schillinger D. Interventions to Improve Care for Patients with Limited Health Literacy. *JCOM*. 2009;16(1):20-29 
 6. Quick Guide to Health Literacy – Fact Sheet. <https://health.gov/communication/literacy/quickguide/factsliteracy.html> (<https://health.gov/communication/literacy/quickguide/factsliteracy.html>). Accessed August 3, 2019 
 7. Koh HK, Brach C, Harris LM, Parchman ML. A Proposed Health Literate Care Model Would Constitute a Systems Approach to Improving Patients Engagement in Care. *Health Aff*. 2013;32(2):357-367. 
 8. JF, Griffith CH, Barnett DR. Residents' Ability to Identify Patients with Poor Literacy Skills. *Acad Med*. 2002;77(10):1039-1041. 
 9. Johnson RL, Saha S, Arbelaez JJ, Beach MC, Cooper LA. Racial and Ethnic Differences in Patient Perceptions of Bias and Cultural Competence in Healthcare. *J Gen Intern Med*. 2004;19(2):101-110 
 10. Baker DW. Health Literacy and Mortality Among Elderly Persons. *Arch Intern Med*. 2007;167(14):1503 
 11. Davis TC, Wolf MS, Bass PF, et al. Low Literacy Impairs Comprehension of Prescription Drug Warning Labels. *J Gen Int Med*. 2006;21(8):847-851. 
 12. Badarudeen S, Sabharwal S. Assessing Readability of Patient Education Materials: Current Role in Orthopaedics. *Clin Orthop Relat Res*. 2010;468(10):2572-2580 
 13. National CLAS Standards – <https://minorityhealth.hhs.gov/omh/browse.aspx?lvl=2&lvlid=53> (<https://minorityhealth.hhs.gov/omh/browse.aspx?lvl=2&lvlid=53>). Accessed October 2, 2017 